

# Variational electrodynamics of atoms

Jayme De Luca\*

Universidade Federal de São Carlos,

Departamento de Física

Rodovia Washington Luis, km 235

Caixa Postal 676, São Carlos, São Paulo 13565-905

(Dated: December 27, 2012)

We study extrema with velocity discontinuities of the *variational* two-body problem with boundary conditions in past and future. Along  $C^2$  segments, these *broken extrema* satisfy Euler-Lagrange equations which are *neutral differential delay* equations with state-dependent deviating arguments. At points where velocities are *discontinuous*, broken extrema (spiky orbits) satisfy the Weierstrass-Erdmann corner conditions that *energies* and *momenta* are *continuous*. We construct *periodic* broken extrema near some  $C^\infty$  circular orbits, using solutions of the neutral differential delay equations along regular segments and a variational approximation for the spiky boundary-layer segments. Finitely measured families of spiky periodic orbits result from the Weierstrass-Erdmann stabilization of the explosive tangent modes near some discrete circular orbits, with scales defined by the Weierstrass-Erdmann conditions. The Weierstrass-Erdmann conditions are *further compatible* with a condition to vanish the far-fields, thus creating a finitely measured family of periodic orbits with vanishing far-fields right at the atomic magnitude. We vary the mass ratio by *three orders of magnitude*, i.e., the *hydrogen*, *muonium* and *positronium* atoms. In each case the broken extrema have magnitudes in quantitative agreement with nature and quantum mechanics, i.e., the spacings between consecutive discrete orbital angular momenta agree with Planck's constant within thirty-percent, while orbital frequencies agree with a corresponding circular line of quantum mechanics within a few percent. The agreement is surprisingly better for the decay from the deepest quantum levels. Last, we discuss the *extension* of classical electrodynamics to *variational* electrodynamics.

Keywords: calculus of variations, neutral differential delay equations, electrodynamics of point charges, state-dependent delay

## INTRODUCTION

The *variational* formulation of classical electrodynamics [1] has continuous extrema with an undefined velocity at some points [2], henceforth *broken extrema*. Along  $C^2$  segments, broken extrema satisfy *neutral differential delay* equations with *state-dependent* deviating arguments [1, 2]. At corner points velocities/accelerations are not defined and broken extrema satisfy the Weierstrass-Erdmann corner conditions (WECC) that *energies* and *momenta* are *continuous* [3, 4]. We study the WECC for the variational electromagnetic two-body problem and its thresholds for corner creation.

We construct periodic broken extrema of variational electrodynamics with a boundary-layer perturbation involving regular segments where the neutral differential delay equations and deviating arguments are *linearized*, while on the (thin) boundary-layer segments a variational approximation is used. Periodic broken extrema (spiky orbits), bifurcate from a discrete set of circular orbits with scales defined by the WECC. We vary the mass-ratio by *three orders of magnitude* to include the atoms of *hydrogen*, *muonium* and *positronium*.

The Weierstrass-Erdmann mechanism of corner creation stabilizes the tangent dynamics of spiky orbits, which are oscillations transversal to the particle separation. In the three cases of hydrogen, muonium and positronium, a discrete set of finitely measured orbits

with vanishing far-fields have frequencies in agreement with quantum mechanics (QM) within a few percent. The qualitative agreement with QM is as follows: (i) the angular momenta of the broken minimizers are approximately integer multiples of a basic angular momentum agreeing with Planck's constant within thirty percent, (ii) the emitted frequency is the difference of two eigenvalues of a suitable linear problem, just like in QM and (iii) the Weierstrass-Erdmann conditions involve precisely the same relevant quantities of QM, i.e., continuity of momenta and energies at corner points.

Maxwell's partial differential equations (PDE) represent the experimentally verified integral laws of classical electrodynamics, e.g., Gauss's surface integral law for the electric field yields the divergence law *when* fields are  $C^1$ . Differential laws were first written by Heaviside sometime after Maxwell completed electrodynamics in the 1860's, and good references exist [5]. Wheeler-Feynman electrodynamics of point charges (1945) [6] was constructed to avoid a shortcoming of Maxwell's PDE's, namely the divergence of the Liénard-Wiechert fields along charge's trajectories. Wheeler and Feynman [6] derived *neutral differential delay equations* of motion (NDDE), whose qualitative behavior was not understood at their time. As natural for the variational principle, solutions of NDDE must be defined piecewise [7, 8]. Continuation of solutions often leaves a set of zero measure where trajectories are *not* differentiable [7, 8].

Quantum mechanics is an enormously successful probabilistic theory [9], and since measure theory is the foundation of probability theory, the interface between variational electrodynamics and QM is here consistently explored with a measure-theoretic generalisation of classical electrodynamics.

Electromagnetic fields of variational electrodynamics are not defined outside trajectories, and must be *extended* with the Liénard-Wiechert formulas. Extrema with discontinuous velocities create *discontinuous* extended fields, often undefined on sets of zero measure. Several results of Maxwell's electrodynamics do not carry over, e.g., Poynting's theorem is valid only in regions where the extended fields are  $C^2$ . Gauss's law can be shown by density arguments to hold for extended fields of trajectories having a *finite* number of corners [2]. Extended fields of trajectories with an *infinite* number of corners per average time unit have not yet been studied. Generalizations of the integral laws of classical electrodynamics using Sobolev's trace theorems [10], and measure-theoretic generalizations to Lebesgue-integrable action functionals, are also open problems.

The electromagnetic two-body problem has a known one-parameter family of  $C^\infty$  solutions, i.e., the circular orbits [11], which create non-vanishing far-fields and are minimizers of variational electrodynamics as well. A surprising result for two-body orbits with repulsive interaction and motion restricted to a line [12] is that globally defined  $C^2$  orbits are uniquely determined by the distance of closest approximation, i.e., a one-parameter invariant manifold. The analogous problem within variational electrodynamics has not been studied yet. The stability of circular orbits was first studied in [13, 14] and [15]. Action-at-a-distance is discussed in [1, 16–18], some electromagnetic-like delay equations are discussed in [19] and the WKB quantization of circular orbits is discussed in [20, 21].

This paper is divided as follows: In Section II we explain variational electrodynamics as an extension of Wheeler-Feynman electrodynamics. We introduce the variational structure, the WECC, and explain the generalized absorber condition needed in variational electrodynamics. In Section III we introduce the Schild circular orbits and magnitudes for later use, which is our starting point to construct a fast-scale (stiff) tangent dynamics. In Section IV we linearize the Wheeler-Feynman NDDE about circular orbits. We explain in detail the infinite-dimensional tangent dynamics of circular orbits and the infinite number of linearly unstable transversal modes. Section V contains the boundary layer theory and the Weierstrass-Erdmann threshold for corner creation. In Section VI we study the magnitudes prescribed by the WECC for a finitely measured family of periodic broken extrema. In Section VI we compare the magnitudes of the broken extrema with nature and QM for *hydrogen*, *muonium* and *positronium*. In Section VII we discuss

the GAC mechanism to vanish the far-fields and show how it is *surprisingly compatible* with the WECC. Last, Section VIII contains discussions, speculations and the conclusion.

## VARIATIONAL ELECTRODYNAMICS

Here we discuss an extension of Wheeler-Feynman's classical electrodynamics of point charges [6]. Unlike Maxwell's electrodynamics, variational electrodynamics does not involve the solution of a PDE: Instead, *finite* segments of charge's trajectories are critical points of a variational principle with Einstein-local boundary conditions in past and future [1, 2]. A  $C^2$  extremum of variational electrodynamics satisfies Euler-Lagrange equations where the Liénard-Wiechert fields of other charges enter as coupling terms. These Euler-Lagrange equations hold only *along* the finite segment of trajectories of the variational problem. Elsewhere in space electromagnetic fields are obtained by *extension*, often modulo a set of zero measure in light-cone with breaking points. Extension of fields to almost everywhere in  $\mathbb{R} \times \mathbb{R}^3$  requires globally defined extrema, otherwise extension is limited inside a ball of finite radius.

We henceforth use a unit system where the speed of light is  $c \equiv 1$  and the electronic and protonic masses and charges are  $m_1 \equiv 1$  and  $q_1 \equiv -1$  and  $m_2 = 1836.1526$  and  $q_2 = 1$ , respectively. Electrodynamics until the 1900's had in mind the Lorentz force equation of motion for charge  $q_i$ , i.e.,

$$m_i \frac{d}{dt} \left( \frac{\mathbf{v}_i}{\sqrt{1 - \mathbf{v}_i^2}} \right) = q_i (\mathbf{E}(\mathbf{x}_i, t) + \mathbf{v}_i \times \mathbf{B}(\mathbf{x}_i, t)), \quad (1)$$

where particle  $i$  has mass  $m_i$  and  $\mathbf{x}_i(t) \in C^2(\mathbb{R})$ . The idea of solving Maxwell's PDE's for the fields  $\mathbf{E}(\mathbf{x}_i, t)$  and  $\mathbf{B}(\mathbf{x}_i, t)$  entered into a crisis when the Liénard-Wiechert general solution for the field of a system of point charges came out [5], i.e.,

$$\mathbf{E}(\mathbf{x}_i, t) = \sum_j (a \mathbf{E}_{j+} + b \mathbf{E}_{j-}), \quad (2)$$

$$\mathbf{B}(\mathbf{x}_i, t) = \sum_j (a \mathbf{B}_{j+} + b \mathbf{B}_{j-}), \quad (3)$$

where summation is over all charges,  $a$  and  $b$  are any pair of real constants satisfying  $a + b = 1$  [5] and

$$\mathbf{E}_{j\pm} \equiv q_j \left\{ \frac{\mathbf{u}_{j\pm}}{\gamma_{j\pm}^2 r_{j\pm}^2} + \frac{\mathbf{n}_{j\pm} \times (\mathbf{u}_{j\pm} \times \mathbf{a}_{j\pm})}{r_{j\pm}} \right\}, \quad (4)$$

$$\mathbf{u}_{j\pm} \equiv \frac{(\mathbf{n}_{j\pm} \pm \mathbf{v}_{j\pm})}{(1 \pm \mathbf{n}_{j\pm} \cdot \mathbf{v}_{j\pm})^3}, \quad (5)$$

$$\mathbf{B}_{j\pm} \equiv \mp \mathbf{n}_{j\pm} \times \mathbf{E}_{j\pm}, \quad (6)$$

with  $\gamma_{j\pm} \equiv (1 - \mathbf{v}_{j\pm}^2)^{-1/2}$ ,  $\mathbf{v}_{j\pm} \equiv d\mathbf{x}_j/dt|_{t=t_{j\pm}}$  and  $\mathbf{a}_{j\pm} \equiv d^2\mathbf{x}_j/dt^2|_{t=t_{j\pm}}$  the velocity and acceleration of charge  $j$

evaluated at the retarded/advanced time  $t_{j\mp}$  defined by

$$t_{j\mp} = t \mp \|\mathbf{x}_j(t_{j\mp}) - \mathbf{x}\|, \quad (7)$$

an implicit state-dependency on  $\mathbf{x}_j(t_{j\mp})$  henceforth called the light-cone condition. Last, in Eqs. (4),  $\mathbf{n}_{j\pm} \equiv (\mathbf{x} - \mathbf{x}_j(t_{j\mp}))/r_{j\pm}$  is a unit vector from position  $\mathbf{x}_j(t_{j\mp})$  to point  $\mathbf{x}$  and  $r_{j\pm} \equiv \|\mathbf{x} - \mathbf{x}_j(t_{j\mp})\|$  denotes distance in light-cone. Maxwell's PDE's fields (4) diverge along each charge's trajectory: Wheeler and Feynman's recipe for a finite Lorentz force [6] was to restrict the summation in Eqs. (2) and (3) to  $j \neq i$ , excluding the divergent self-field.

The choice  $i \neq j$  combined with  $a = b = 1/2$  is the solution of the variational boundary value problem explained bellow. Even restricted to almost everywhere subluminal, non-collisional trajectories[1], the electromagnetic action principle is more complex than the two-point boundary value principle of Newtonian mechanics since boundary segments in past and future must be given to provide data for the advanced and retarded arguments [1], as illustrated in FIG.(1). The needed Einstein-local boundary conditions consist of (a) the initial point  $O_1$  for trajectory 1 and respective boundary-segment of trajectory 2 inside the light-cone of  $O_1$ , and (b) the final point  $L_2$  for trajectory 2 and respective boundary-segment of trajectory 1 inside the light-cone of  $L_2$ . The variational problem is to find the segments of trajectories between the endpoints  $O_1$  and  $L_2^-$ , and  $O_1^+$  and  $L_2$  of FIG.(1). As shown in [18] and illustrated in FIG. (1), along almost everywhere subluminal continuous orbits satisfying these boundaries, the light-cone conditions (7) define unique solutions for the deviating arguments  $t_{j\pm}(t, \mathbf{x}_i(t))$ .

The action functional is a sum of four integrals: two *local* integrals each involving one trajectory, i.e.,  $\int T_i(\mathbf{x}_i, \dot{\mathbf{x}}_i) dt_i$ , and two *interaction* integrals of functions of both positions/velocities, where one position/velocity is evaluated at a deviating time argument,  $\int V_{ij}^\pm(\mathbf{x}_i, \dot{\mathbf{x}}_i, \mathbf{x}_{j\pm}, \dot{\mathbf{x}}_{j\pm}) dt_i$ . The action can be expressed in two equivalent forms, i.e.,

$$S \equiv \int_{t_{O_1^+}}^{t_{L_2}} T_2 dt_2 + \int_{t_{O_1}}^{t_{L_2^-}} T_1 dt_1 + \underbrace{\int_{t_{O_1}}^{t_{L_2^+}} V_{12}^- dt_1}_{\parallel} + \underbrace{\int_{t_{O_1}}^{t_{L_2^-}} V_{12}^+ dt_1}_{\parallel}, \quad (8)$$

$$= \int_{t_{O_1}}^{t_{L_2^-}} T_1 dt_1 + \int_{t_{O_1^+}}^{t_{L_2}} T_2 dt_2 + \underbrace{\int_{t_{O_1^-}}^{t_{L_2}} V_{21}^+ dt_2}_{\parallel} + \underbrace{\int_{t_{O_1^+}}^{t_{L_2^-}} V_{21}^- dt_2}_{\parallel}, \quad (9)$$

where vertical braces indicate equivalence by change of the integration variable on the interaction integrals. The Jacobian for the change of variables is the derivative of the delayed times  $t_{j\pm}(t_i, \mathbf{x}_j(t_i))$  evaluated along the orbit, as obtained taking a derivative of the implicit condi-




FIG. 1. Schematic showing the Einstein-local boundaries in  $\mathbb{R}^4$ , i.e., (a) initial point  $O_1 \equiv (t_{O_1}, \mathbf{x}_1(t_{O_1}))$  of trajectory 1 and boundary segment  $\mathbf{x}_2(t_2)$  for  $t_2 \in [t_{O_1^-}, t_{O_1^+}]$  inside the light-cone of  $O_1$ , and (b) endpoint  $L_2 \equiv (t_{L_2}, \mathbf{x}_2(t_{L_2}))$  of trajectory 2 and boundary segment  $\mathbf{x}_1(t_1)$  for  $t_1 \in [t_{L_2^-}, t_{L_2^+}]$  inside the light-cone of  $L_2$ . Trajectories  $\mathbf{x}_1(t_1)$  for  $t_1 \in [t_{O_1}, t_{L_2^-}]$  (green line) and  $\mathbf{x}_2(t_2)$  for  $t_2 \in [t_{O_1^+}, t_{L_2}]$  (blue line) are determined by the extremum condition. Black dotted lines indicate the light-cone relation and brown dotted lines indicate the  $V_{12}^-$  and  $V_{12}^+$  interactions. Arbitrary units.

tion (7) with  $\mathbf{x} = \mathbf{x}_i(t_i)$ ,

$$\frac{dt_{j\pm}}{dt_i} = \frac{(1 \pm \mathbf{n}_{j\pm} \cdot \mathbf{v}_i)}{(1 \pm \mathbf{n}_{j\pm} \cdot \mathbf{v}_{j\pm})}, \quad (10)$$

and explained in Refs. [1, 18]. One can thus express the interaction terms by either integrals over  $t_1$  (as in (8)) or integrals over  $t_2$  (as in (9)). A generic variational structure can be defined using arbitrary  $V$ 's on line (8), while the  $V$ 's on line (9) are determined by changing variables with (10) or the equivalent Jacobian if the constraint is other than (7). Henceforth we concentrate on variational electrodynamics, i.e.,

$$T_i \equiv m_i[1 - \sqrt{1 - \mathbf{v}_i^2}], \quad (11)$$

$$V_{ij}^\pm(\mathbf{x}_i, \dot{\mathbf{x}}_i, \mathbf{x}_{j\pm}, \dot{\mathbf{x}}_{j\pm}) \equiv \frac{(1 - \mathbf{v}_i \cdot \mathbf{v}_{j\pm})}{2r_{j\pm}(1 \pm \mathbf{n}_{j\pm} \cdot \mathbf{v}_{j\pm})}, \quad (12)$$

where  $i = 1, 2$  and  $j = 3 - i$ .

For the linear variation, trajectory 2 is to be kept fixed while trajectory 1 is varied, and line (8) with the first term constant defines partial Lagrangian 1. Vice-versa, trajectory 1 is to be kept fixed while trajectory 2 is varied, and line (9) with the first term frozen defines partial Lagrangian 2. A  $C^2$  extremum satisfies the Euler-Lagrange equation of each partial Lagrangian [1], i.e., Eq.(1), the NDDE with state-dependent deviating arguments of Wheeler and Feynman[6].

Next in complexity are *broken extrema* with a finite number of corner points [4]. For each partial Lagrangian,

broken extrema satisfy the WECC that *momenta* and *energies* are *continuous* at corner points. The *first Weierstrass-Erdmann corner condition*, (WECC1), is the continuity of the momentum of partial Lagrangian  $i$ , i.e.,

$$\frac{\partial \mathcal{L}_i}{\partial \mathbf{v}_i} = \frac{m_i \mathbf{v}_i}{\sqrt{1 - \mathbf{v}_i^2}} - \frac{\mathbf{v}_{j-}}{2r_{j-}(1 - \mathbf{n}_{j-} \cdot \mathbf{v}_{j-})} - \frac{\mathbf{v}_{j+}}{2r_{j+}(1 + \mathbf{n}_{j+} \cdot \mathbf{v}_{j+})}, \quad (13)$$

$$\equiv \frac{m_i \mathbf{v}_i}{\sqrt{1 - \mathbf{v}_i^2}} - \mathbf{A}_j, \quad (14)$$

where  $\mathcal{L}_1$  is the sum of the last three integrands of line (8),  $\mathcal{L}_2$  the sum of the last three integrands of line (9) and  $\mathbf{A}_j$  the vector potential of particle  $j$ . Equation (13) holds for  $i = 1, 2$  despite any velocity discontinuities of charge  $j = 3 - i$ . If the extremum has a corner for particle  $i$  at time  $t_i$ , the trajectory of particle  $j$  compensates with a corner in light-cone at either  $t_{j-}$  or  $t_{j+}$ .

For the *second Weierstrass-Erdmann corner condition*, (WECC2), we express the partial Lagrangian  $i$  as

$$\mathcal{L}_i = T_i - \mathbf{v}_i \cdot \mathbf{A}_j + U_j, \quad (15)$$

where

$$\mathbf{A}_j \equiv \frac{\mathbf{v}_{j-}}{2r_{j-}(1 - \mathbf{n}_{j-} \cdot \mathbf{v}_{j-})} + \frac{\mathbf{v}_{j+}}{2r_{j+}(1 + \mathbf{n}_{j+} \cdot \mathbf{v}_{j+})},$$

$$U_j \equiv \frac{1}{2r_{j-}(1 - \mathbf{n}_{j-} \cdot \mathbf{v}_{j-})} + \frac{1}{2r_{j+}(1 + \mathbf{n}_{j+} \cdot \mathbf{v}_{j+})}.$$

The *partial energy* of partial Lagrangian  $i$  evaluates to

$$E_i \equiv \mathbf{v}_i \cdot \frac{\partial \mathcal{L}_i}{\partial \mathbf{v}_i} - \mathcal{L}_i = \frac{m_i}{\sqrt{1 - \mathbf{v}_i^2}}. \quad (16)$$

Therefore, the WECC2 is simply that  $|\mathbf{v}_i|$  must be *continuous* at each breaking point of each trajectory, in order to make the partial energies  $E_i$  continuous.

Next we express the WECC1 using an upper index  $l$  or  $r$  to indicate respectively left-velocity or right-velocity at the breaking point. Using (13) and the WECC2, Eq. (16), the WECC1 becomes

$$\Delta\left(\frac{\partial \mathcal{L}_i}{\partial \mathbf{v}_i}\right) = \frac{m_i \Delta \mathbf{v}_i}{\sqrt{1 - \mathbf{v}_i^2}} - \left( \frac{\Delta \mathbf{v}_{j-} - \mathbf{n}_{j-} \times (\mathbf{v}_{j-}^r \times \mathbf{v}_{j-}^l)}{2r_{j-}(1 - \mathbf{n}_{j-} \cdot \mathbf{v}_{j-}^l)(1 - \mathbf{n}_{j-} \cdot \mathbf{v}_{j-}^r)} \right) - \left( \frac{\Delta \mathbf{v}_{j+} + \mathbf{n}_{j+} \times (\mathbf{v}_{j+}^r \times \mathbf{v}_{j+}^l)}{2r_{j+}(1 + \mathbf{n}_{j+} \cdot \mathbf{v}_{j+}^l)(1 + \mathbf{n}_{j+} \cdot \mathbf{v}_{j+}^r)} \right) \equiv 0, \quad (17)$$

where  $\Delta \mathbf{v}_i \equiv \mathbf{v}_i^r - \mathbf{v}_i^l$ ,  $\Delta \mathbf{v}_{j\pm} \equiv \mathbf{v}_{j\pm}^r - \mathbf{v}_{j\pm}^l$ .

Last, we already mentioned that fields of trajectories defined in  $t \in [-\infty, \infty]$ , henceforth *globally bounded*, can be *extended* to almost everywhere in  $\mathbb{R} \times \mathbb{R}^3$ . By *extension* we mean fields defined by formulas (4) and

(6) *whenever* all the respective retarded/advanced velocities/accelerations are well defined. At least for trajectories with a finite number of corners per unit time, formulas (4) and (6) define extended fields almost everywhere but on a set of zero measure. Extension is used when another charge is added to the system, at which stage minimization respect to its trajectory yields an extra Euler-Lagrange equation involving the extended fields. Let us first assume globally bounded trajectories  $\in C^2(\mathbb{R})$ , which implies that all advanced and retarded normals coincide at a far distance, i.e.,  $\mathbf{n}_{j+} = \mathbf{n}_{j-} \equiv \mathbf{n}$ ,  $\forall j$  [2], yielding the extended electric field

$$\mathbf{E}(\mathbf{x}, t) = \frac{1}{2} \sum_j (\mathbf{E}_{j+}(\mathbf{x}, t) + \mathbf{E}_{j-}(\mathbf{x}, t)), \quad (18)$$

while Eq. (6) with  $\mathbf{n}_{j+} = \mathbf{n}_{j-} \equiv \mathbf{n}$  yields

$$\mathbf{B}(\mathbf{x}, t) = \frac{1}{2} \mathbf{n} \times \sum_j (\mathbf{E}_{j-}(\mathbf{x}, t) - \mathbf{E}_{j+}(\mathbf{x}, t)). \quad (19)$$

According to Eq. (1), for an extra distant charge to travel undisturbed at a far distance with an *arbitrary* velocity  $\mathbf{v}_i$ , the *semi-sum* (18) and *semi-difference* (19) of universal electric fields must vanish asymptotically. Wheeler and Feynman's *absorber condition* was obtained using Maxwell's PDE's [6, 22], while the above pre-minimizer condition applies also to piecewise  $C^2$  globally bounded extrema, yielding the separate vanishing *almost everywhere* of the retarded and advanced universal far fields. We henceforth call this the *generalized absorber condition* (GAC) of variational electrodynamics.

Notice that functional (8) is semi-bounded ( $S \geq 0$ ), and under some compactness properties one extremum should be a *minimum* [23]. It is an open question if (8) could also have saddle-point type extrema [18].

In physics, the name *classical trajectory* is synonymous of global solution of an ordinary differential equation (ODE), i.e., a set of points invariant by time evolution under the ODE. In variational electrodynamics, the analogous concept is a global  $C^0$  curve out of which one can arbitrarily cut *finite boundary segments*, as defined above FIG. 1. The invariant manifold can have corners and it is only necessary that *one* extremum of each finite variational problem be a segment of the manifold. Unlike the case of ODE's, one can not continue an invariant manifold forwards (nor backwards) with a time integration. Instead, the infinite dimensional problem at hand is finding a whole function, just like solving a PDE with boundary conditions.

Last and again, trajectories with derivative discontinuities obviously do not have a Taylor series approximation. Together with the failure of Poynting's theorem for extended fields of spiky orbits, these are two major generalisations of Dirac's classical electrodynamics of point charges [24]. The generalization is welcome after

the proof of Ref. [2] that globally bounded two-body orbits with vanishing far-fields *must* have corners.

### CIRCULAR ORBITS

The details of the  $C^\infty$  Schild circular orbits [11] are given below using the notation of Ref. [15]: The constant angular velocity and distance in light-cone are denoted by  $\Omega$  and  $r_b$ , respectively, and the angle  $\theta \equiv \Omega r_b$  that particles turn in the light-cone time is henceforth called the *delay angle*. The family of subluminal  $C^\infty$  circular orbits [11] is parametrized by  $\theta \in (0, 1)$ , and for atomic orbits  $\theta \lesssim 10^{-2}$ . Along orbits with a small delay angle  $\theta$ , the Kepler formulas yield the leading order angular velocity and distance in light-cone, respectively, i.e.,

$$\Omega = \mu\theta^3 + O(\theta^5), \quad (20)$$

$$r_b = \frac{\theta}{\Omega} = \frac{1}{\mu\theta^2}. \quad (21)$$

where reduced mass and total mass are defined by  $\mu \equiv m_1 m_2 / M$  and  $M \equiv m_1 + m_2$ . It is important to keep these dependencies in mind, and that for hydrogen  $\mu \simeq (1836/1837) \simeq 1$ . Adopting the same notation of Ref. [15], we express each particle's circular orbit's radius by numbers  $0 \leq b_i < 1$  as

$$r_i \equiv b_i r_b, \quad (22)$$

which define scalar velocities

$$v_i = \Omega r_i = \theta b_i, \quad (23)$$

for  $i = 1, 2$ .

As illustrated in FIG. (2) and in Ref. [11], the circular radii and the distance in light-cone form a triangle of largest side  $r_b$ , yielding a trigonometric constraint

$$b_1^2 + b_2^2 + 2b_1 b_2 \cos(\theta) = 1, \quad (24)$$

equivalent to Eq. (3.1) of Ref. [11]. Using the ratio of equations (3.2) and (3.3) of Ref. [11] to eliminate  $\Omega$ , together with constraint (24), we find the solution  $b_1 = (1 + \frac{\mu\theta^2}{2M}) \frac{m_2}{M} + O(\theta^4)$  and  $b_2 = (1 + \frac{\mu\theta^2}{2M}) \frac{m_1}{M} + O(\theta^4)$ .

Last, Noether's angular momentum

$$l_z = \frac{1 + b_1 b_2 \theta^2 \cos(\theta)}{\theta + b_1 b_2 \theta^2 \sin(\theta)} \simeq \frac{1}{\theta}, \quad (25)$$

is an important quantity of the circular orbit to keep in mind [11, 14, 20]. Atomic orbits have  $l_z$  of the order of one over the fine-structure constant,  $\hbar c / e^2 = 137.036$ , a fundamental magnitude of nature. Delay angles and angular momenta of Bohr orbits in our unit system are respectively  $\theta \simeq (e^2 / \hbar c) / q$  and  $l_z \simeq (\hbar c / e^2) q$ , for each nonzero integer  $q$ .




FIG. 2. Schild circular orbit with particles in diametral opposition at the same time of the inertial frame (dotted straight line is  $t = 0$ ). Indicated is the retarded position of particle 1 along the outer circle, and the angle  $\theta$  travelled during the light-cone time. Particle separation ( $r_b$ ) in the past light-cone of particle 2 is the red line making an angle  $\psi_2$  with the dotted line. The delay angle and the protonic circular radius have an exaggerated magnitude for illustrative purposes. Arbitrary units.

### LINEAR MODES

In order to linearize the equations of motion about circular orbits of small delay angles, we re-write the Lorentz-force Eqs. of motion (1) as

$$\frac{m_i \mathbf{a}_i}{\sqrt{1 - \mathbf{v}_i^2}} = q_i [\mathbf{E}_j - (\mathbf{v}_i \cdot \mathbf{E}_j) \mathbf{v}_i + \mathbf{v}_i \times \mathbf{B}_j], \quad (26)$$

by evaluating the derivative on the left-hand side of Eq. (1) and subtracting the scalar product with  $\mathbf{v}_i$ , where  $i = 1, 2$  and  $j = 3 - i$ . The magnetic term (last term on the right-hand side of (26)) is a transversal force proportional to the electric field by (6), and further proportional to the velocity, which is small for small  $\theta$  (see Eq. (23)).

The electric force, first term on the right-hand side of Eq. (26), is responsible for the significant contributions to the linearized equations along a circular orbit: The electric field (4) decomposes in two terms: (i) the near-electric field proportional to  $1/r_{j\pm}^2$ , a magnitude of  $\mu^2 \theta^4$  by use of (21), times an (almost) radial vector, and (ii) the far-electric field proportional to  $1/r_{j\pm}$ , a magnitude of  $\mu \theta^2$  by use of (21), times a vector (almost) along the circular orbit. It is important to ponder upon the magnitude of each electric contribution: the far-electric field is (almost) transversal, and projection to the normal direction brings another factor of  $\theta$ . The force along circular orbits is approximately described by the near field only *because* the far-field (4) is further proportional to the acceleration (which is  $O(1/r^2)$  along circular orbits). The contributions to the right-hand-side of (26) have magni-

tudes

$$F_i \propto O(\theta^4) + |\mathbf{a}_j|O(\theta^3). \quad (27)$$

In Eq. (27), the near-field contribution is  $O(\theta^4)$  while the far-field contribution is  $|\mathbf{a}_j|O(\theta^3)$ , which is smaller along circular orbits since  $|\mathbf{a}_j| \propto 1/r^2 = O(\theta^4)$ . Upon linearization about the circular orbit, this dominance changes: the linearized equations accept solutions of arbitrarily large frequencies, and the most important linearized term is precisely the *far-field term*.

Next we derive the linearized equations along the orbital plane using the notation of Ref. [15]. We introduce complex gyroscopic coordinates where the circular orbit is a fixed point of the equations of motion, i.e.,

$$x_j + iy_j \equiv r_b \exp(-i\Omega t)[b_j + l_j + iu_j], \quad (28)$$

where  $(l_j, u_j)$  are respectively the longitudinal and transversal gyroscopic coordinates. The circular orbit [11] is the fixed point  $(l_j, u_j) = (0, 0)$  for  $j = 1, 2$ . The dominant linear correction for large accelerations (the stiff limit [15]) is obtained from (26) with only the first term on the right-hand side and far-field  $\mathbf{E}_{j\pm}$  approximated by

$$\mathbf{E}_{j\pm} \simeq \frac{1}{r_b} \mathbf{n}_{j\pm} \times (\mathbf{n}_{j\pm} \times \delta \mathbf{a}_{j\pm}) = \frac{1}{r_b} (\mathbf{n}_{j\pm} \cdot \delta \mathbf{a}_{j\pm}) \mathbf{n}_{j\pm} - \frac{1}{r_b} \delta \mathbf{a}_{j\pm}, \quad (29)$$

where  $\pm$  indicate evaluation at the unperturbed deviating times  $t \pm (\theta/\Omega)$  (because we want the linear term only). The gyroscopic representation of the rotating normals in light-cone are complex numbers of unit modulus, i.e.,  $\exp(-i\Omega t) \exp(i\psi_j)$ , where

$$\begin{aligned} \exp(i\psi_1) &\equiv b_1 + b_2 \exp i\theta, \\ \exp(i\psi_2) &\equiv b_2 + b_1 \exp i\theta. \end{aligned} \quad (30)$$

The unitarity of (30) is seen using Eq. (24). Angles  $\psi_1$  and  $\psi_2$  further satisfy  $\psi_1 + \psi_2 = \theta$ , and for small  $\theta$  the  $b_i$  given below Eq. (24) yield  $\psi_1 \simeq \frac{m_1}{M}\theta$  and  $\psi_2 \simeq \frac{m_2}{M}\theta$ . In FIG. (2),  $\psi_2$  is the angle between the (rotating) red line and the  $x$ -axis (dashed line).

The linearized planar equations of motion are obtained substituting (28) into (26) with only its first right-hand-side term given by the semi-sum of retarded and advanced fields (29). The real and complex parts of the linearized equations of motion keeping only the largest derivatives of the gyroscopic coordinates yields

$$\begin{aligned} m_1 r_b \ddot{l}_1 &= -C_{11} \frac{(\ddot{l}_{2+} + \ddot{l}_{2-})}{2} - S_{11} \frac{(\ddot{u}_{2+} - \ddot{u}_{2-})}{2}, \\ m_2 r_b \ddot{l}_2 &= -C_{12} \frac{(\ddot{l}_{1+} + \ddot{l}_{1-})}{2} - S_{12} \frac{(\ddot{u}_{1+} - \ddot{u}_{1-})}{2}, \\ m_1 r_b \ddot{u}_1 &= -C_{13} \frac{(\ddot{u}_{2+} + \ddot{u}_{2-})}{2} - S_{13} \frac{(\ddot{l}_{2+} - \ddot{l}_{2-})}{2}, \\ m_2 r_b \ddot{u}_2 &= -C_{14} \frac{(\ddot{u}_{1+} + \ddot{u}_{1-})}{2} - S_{14} \frac{(\ddot{l}_{1+} - \ddot{l}_{1-})}{2}, \end{aligned} \quad (31)$$

where  $C_{1j}$  and  $S_{1j}$  are given below Eq. (33). Equation (31) is a *linear* NDDE with exponential solutions

$$\begin{aligned} l_j &= L_j \exp(\lambda \Omega t / \theta), \\ u_j &= U_j \exp(\lambda \Omega t / \theta), \end{aligned} \quad (32)$$

where  $i, j = 1, 2$  and  $L_1, L_2, U_1, U_2$  are non-trivial solutions of

$$\begin{pmatrix} m_1 r_b & D_{11} & 0 & E_{11} \\ D_{12} & m_2 r_b & E_{12} & 0 \\ 0 & E_{13} & m_1 r_b & D_{13} \\ E_{14} & 0 & D_{14} & m_2 r_b \end{pmatrix} \begin{pmatrix} L_1 \\ L_2 \\ U_1 \\ U_2 \end{pmatrix} = 0, \quad (33)$$

where

$$D \equiv (\cosh \lambda) C = -\cosh \lambda \begin{pmatrix} \cos \theta - \cos \psi_2 \cos \theta \\ \cos \theta - \cos \psi_1 \cos \theta \\ \cos \theta + \sin \psi_2 \sin \theta \\ \cos \theta + \sin \psi_1 \sin \theta \end{pmatrix}, \quad (34)$$

and

$$E \equiv (\sinh \lambda) S \equiv -\sinh \lambda \begin{pmatrix} \sin \theta - \sin \psi_2 \cos \theta \\ \sin \theta - \sin \psi_1 \cos \theta \\ (1 - \cos \psi_2) \sin \theta \\ (1 - \cos \psi_1) \sin \theta \end{pmatrix}. \quad (35)$$

A nontrivial solution of (33) requires the vanishing of its  $4 \times 4$  determinant,

$$F_{xy} = 1 - \frac{\mu \theta^4}{M} \cosh^2(\lambda) = 0, \quad (36)$$

where powers of  $\theta^2 \cosh \lambda$  with coefficients of  $O(\theta^4)$  were discarded, henceforth called the stiff-limit [15]. The roots of Eq. (36) exist in symplectic sets of four, i.e.,  $(\lambda, -\lambda, \lambda^*, -\lambda^*)$ . For atomic hydrogen,  $\theta^{-1} \sim 137.036$  and  $(\mu/M) \simeq (1/1836)$ , so that Eq. (36) requires  $\lambda$  to have a real part  $|\Re(\lambda)| \equiv \sigma = \ln(\sqrt{\frac{4M}{\mu\theta^4}}) \simeq 14.29$ , while the imaginary part of  $\lambda$  is an integer multiple of  $\pi i$ . The general solution of Eq. (36) modulo the symplectic symmetry is

$$\lambda = \sigma + \pi q i, \quad (37)$$

where  $i \equiv \sqrt{-1}$  and  $q \in \mathbb{Z}$ . Solution (37) was called a ping-pong mode in Ref. [15] because its phase advances by  $\pi q$  in one light-cone time  $r_b$ , i.e., a phase speed of  $\pi q \Omega / \theta = \mu \theta^2 \pi q = \pi q / r_b = (\Omega / \theta) \Im(\lambda)$ . The only  $O(1)$  off-diagonal terms of matrix (33) are  $D_{13}$  and  $D_{14}$ , others being  $O(\theta)$  or higher order. The nontrivial eigenvector solution is approximately

$$(L_1, L_2, U_1, U_2) \propto \left( \frac{\mu \theta}{M}, \frac{\mu \theta}{M}, 1, \sqrt{\frac{\mu}{M}} \right), \quad (38)$$

so that for  $m_2 \gg m_1$  solution (32) is along the circular orbit, since  $U_i \gg L_j$ , i.e., a quasi-*transversal* mode. The largest longitudinal component is for positronium at moderate  $\theta$ , but still small. Solution (37) defines a nonzero real part for  $\lambda$ , so that amplitudes blow up at either  $t \rightarrow \pm\infty$ , implying that besides circular orbits there are no other  $C^\infty$  almost-circular orbits defined for all times (i.e., globally bounded).

The inclusion of  $O(1/\lambda)$  and  $O(1/\lambda^2)$  linear terms to the planar motion is outlined in Ref. [15]. The linearized motion perpendicular to the orbital plane is studied analogously. As explained in [15], the  $z$ -oscillations are transversal modes that decouple from the planar transversal oscillations (31) at linear order. The determinant of the linearized  $2 \times 2$  system is again (36), an *asymptotic* degeneracy. The degeneracy is raised by  $O(1/\lambda^2)$  corrections to (33) and (36) introduced by the linear terms with lower derivatives [15]. The determinant for planar modes is calculated in Ref. [15] up to  $O(\frac{1}{\lambda^4})$  terms (see Eq. (41) of Ref. [15] with  $\Gamma = -1/2$ ), i.e.,

$$F_{xy} = 1 - \frac{\mu\theta^4}{M} \left(1 + \frac{7}{\lambda^2} + \frac{5}{\lambda^4}\right) \cosh^2(\lambda) + \frac{\mu\theta^4}{M} \left(\frac{1}{\lambda} + \frac{5}{\lambda^3}\right) \sinh(2\lambda) = 0, \quad (39)$$

while the determinant for perpendicular oscillations and up to  $O(\frac{1}{\lambda^4})$  terms is Eq. (B17) in appendix B of [15] with  $\Gamma = -1/2$ , i.e.,

$$F_z = 1 - \frac{\mu\theta^4}{M} \left(1 - \frac{1}{\lambda^2} + \frac{1}{\lambda^4}\right) \cosh^2(\lambda) + \frac{\mu\theta^4}{M} \left(\frac{1}{\lambda} - \frac{1}{\lambda^3}\right) \sinh(2\lambda) = 0. \quad (40)$$

Notice the symplectic symmetry that roots of Eqs. (39) and (40) are still in sets of four,  $(\lambda, -\lambda, \lambda^*, -\lambda^*)$ , and even the first correction at  $O(\frac{1}{\lambda})$  is the same. The corrections at  $O(\frac{1}{\lambda^2})$  separate Eq. (39) from (40), unfolding the asymptotic degeneracy of planar and perpendicular transversal modes.

### BOUNDARY LAYER

The non-zero real part in Eq. (37) suggests that there are no other near-circular  $C^\infty$  solutions besides circular orbits, since all linearized modes blow up at either  $t \rightarrow \pm\infty$ . This unless a corner point invalidates the linearization. Moreover, Ref. [2] already found that globally bounded two-body orbits with vanishing far-fields (the GAC) *must* have velocity discontinuities. Therefore, we are led to seek globally bounded extrema *with corners*.

We set about the task to construct a periodic broken extremum using a boundary-layer perturbation that assumes a number of piecewise  $C^\infty$  layers separated by boundary layers each containing one or more corner points, as illustrated in FIG. (3). Along both trajectories, boundary layers have an angular width  $\alpha\theta \ll \theta$ . Outside boundary layers we can linearize because trajectories are  $C^\infty$  and deviating arguments fall in  $C^\infty$  segments. Inside boundary layers we *do not* linearize but rather use a variational approximation for the minimizer with corners and apply the WECC.




FIG. 3. Unperturbed circular orbits (dashed lines), boundary layers along orbit of (heavier) particle 1 (solid red) diving to a corner at a smaller inner radius (dotted line). Boundary layers along orbit 2 (solid blue) spring up to corners at a larger outer radius (dotted line). At points (h), (a) and (c), perturbed orbit 1 (dark solid line) crosses the unperturbed orbit and makes a regular segment with an exaggerated outer radius. At points (d) and (f), perturbed orbit 2 (inner dark circular line) crosses the unperturbed orbit and makes a regular segment at a smaller radius. Boundary layer magnitudes are exaggerated for illustrative purposes. Arbitrary units.

The exact roots of Eqs. (39) and (40) near the limiting root (37) are, respectively,

$$\lambda_{xy}(\theta, q) \equiv \sigma_{xy} + \pi qi + i\epsilon_{xy}, \quad (41)$$

$$\lambda_z(\theta, q) \equiv \sigma_z + \pi qi + i\epsilon_z, \quad (42)$$

where  $q$  is an arbitrary integer,  $\epsilon_{xy}(\theta, q) = -\epsilon_{xy}(\theta, -q)$  and  $\epsilon_z(\theta, q) = -\epsilon_z(\theta, -q)$  (by the symplectic symmetry). An  $O(\mu\theta^4/M)$  approximation for positive  $\sigma$  is obtained by setting  $4\cosh^2(\lambda) \approx 2\sinh(2\lambda) \approx \exp(2\lambda)$  into Eqs. (39) and (40), thus defining polynomials

$$F_{xy}(\lambda) = 1 - \frac{\mu\theta^4}{4M} \exp(2\lambda) \left(1 - \frac{2}{\lambda} + \frac{7}{\lambda^2} - \frac{10}{\lambda^3} + \frac{5}{\lambda^4}\right) \equiv 1 - \frac{\mu\theta^4}{4M} \exp(2\lambda) p_{xy}\left(\frac{1}{\lambda}\right) = 0, \quad (43)$$

and

$$F_z(\lambda) = 1 - \frac{\mu\theta^4}{4M} \exp(2\lambda) \left(1 - \frac{2}{\lambda} - \frac{1}{\lambda^2} + \frac{2}{\lambda^3} + \frac{1}{\lambda^4}\right) \equiv 1 - \frac{\mu\theta^4}{4M} \exp(2\lambda) p_z\left(\frac{1}{\lambda}\right) = 0. \quad (44)$$

It follows from Eqs. (39) and (41) that

$$\exp(4i\epsilon_{xy}) = \frac{p_{xy}(\frac{1}{\lambda_{xy}^*})}{p_{xy}(\frac{1}{\lambda_{xy}})}, \quad (45)$$

while Eqs. (40) and (42) yield

$$\exp(4i\epsilon_z) = \frac{p_z(\frac{1}{\lambda_z^*})}{p_z(\frac{1}{\lambda_z})}. \quad (46)$$

Disregarding the contribution of  $\epsilon_{xy}$  and  $\epsilon_z$  and replacing  $\lambda_{xy} = \lambda_z = \sigma + \pi qi$  in the right-hand-sides of (45) and (46) yields, to the first order in  $(1/\lambda)$

$$\epsilon_{xy} \simeq \epsilon_z = \Im\left(\frac{1}{\lambda}\right) = \frac{-\pi q}{(\sigma^2 + \pi^2 q^2)} \quad (47)$$

We henceforth use a positive integer  $q$ , so that the *energetic mismatches*  $\epsilon_{xy}$  and  $\epsilon_z$  are negative  $O(1/\lambda)$  numbers. Another order can be gained expanding the right-hand-side of (45) in a Taylor series on the deviation  $\epsilon_{xy}$  about  $\lambda_q = \sigma + \pi qi$ , which generates only terms at  $O(1/\lambda^3)$ , so that up to  $O(1/\lambda^2)$  Eq. (45) yields

$$\epsilon_{xy}(q) = \frac{-\pi q}{(\sigma^2 + \pi^2 q^2)} + \frac{5\pi q \sigma}{(\sigma^2 + \pi^2 q^2)^2}, \quad (48)$$

$\epsilon_{xy}$  negative and monotonically increasing for  $q \geq 6$ . Analogously, the right-hand-side of (46) evaluated at  $\lambda_q = \sigma + \pi qi$  yields  $\epsilon_z$  up to  $O(1/\lambda^2)$

$$\epsilon_z(q) = \frac{-\pi q}{(\sigma^2 + \pi^2 q^2)} - \frac{3\pi q \sigma}{(\sigma^2 + \pi^2 q^2)^2}, \quad (49)$$

again negative and monotonically increasing for  $q \geq 6$ .

Let us assume a corner at  $t = 0$  along orbit 1, as illustrated in FIG. (3). We define the first regular layer  $\alpha r_b < t < r_b - \alpha r_b$  as the first and last zeros of the (exponentially increasing) perturbation inside the first light-cone time zone. This is illustrated in FIG. (3) as points (a) and (h) where the perturbed trajectories (solid lines) cross the unperturbed orbits (dotted lines). From (a) to (h) the velocity of the perturbed orbit changes almost by  $\pi q$ . Using a linear combination of the symplectic quartet of linearized modes, and (28) and (38) with  $l_1 \propto L_1 = \frac{\mu\theta}{M}$  and  $u_1 \propto U_1 = 1$ , the orbital perturbation constructed to vanish at layer edges is

$$\begin{pmatrix} \Delta x_j \\ \Delta y_j \end{pmatrix} = \begin{pmatrix} \frac{\mu\theta A}{M} \\ A \end{pmatrix} \frac{\cosh\left(\frac{\sigma_{xy} t}{r_b}\right)}{\cosh\left(\frac{\sigma_{xy}}{2}\right)} \sin\left(\frac{[\pi q + \epsilon_{xy}(q) - \theta](t - \alpha r_b)}{r_b}\right). \quad (50)$$

In Eq. (50),  $\sigma_{xy}$  and  $\epsilon_{xy}$  are given by (41) and we must have the amplitude  $A < \frac{r_b}{\pi q}$ , to avoid a superluminal velocity at layer edges. Along the regular region the phase of the sine function in (50) advances by almost  $\pi q$ , and the condition of vanishing at  $t = r_b - \alpha r_b$  yields  $\alpha = \frac{(\epsilon_{xy} - \theta)}{2\pi q}$ .

An instructive variational approximation for the boundary layer motion is illustrated in FIG. (3), with corners along a given orbit all equivalent by a  $\theta$ -rotation. The central spike starts along trajectory 1 with a ninety-degree corner, point (a), and respective corner in light-cone along trajectory 2, point (d). Next is a straight-line dive ( $\bar{ab}$ ) to an almost 180-degree corner ((b) and (e)), then a straight-line climb ( $\bar{bc}$ ) to the last ninety-degree

corner ((c) and (f)), resuming motion along the next regular layer.

Corners satisfy the WECC, and 180-degree corners, (points (a) and (d) in FIG. (3)), are simpler to analyze: For planar motion, condition (17) yields four equations, one along each Cartesian direction and for  $i = 1, 2$ . Assuming 180-degree corners are reflections about the light-cone direction  $\mathbf{n}$ , the vectorial components of (17) perpendicular to each  $\mathbf{n}$  vanish trivially by symmetry. The two nontrivial conditions yield a  $2 \times 2$  linear homogeneous system for the  $\mathbf{n} \cdot \Delta \mathbf{v}_j$ , with determinant

$$D = m_1 m_2 \sqrt{1 - v_1^2} \sqrt{1 - v_2^2} - \frac{1}{\bar{r}^2}, \quad (51)$$

where  $\bar{r}$  is distance in light-cone from (a) to (d). In deriving (51) from (17) we assumed 180-degrees reflections, otherwise there is a contribution from the  $(\mathbf{v}_{j\pm}^r \times \mathbf{v}_{j\pm}^l)$  terms. Ninety degree corners are analyzed substituting  $\mathbf{v}_i^r = \mathbf{v}_i^l + \Delta \mathbf{v}_i$  into (17), yielding a  $4 \times 4$  linear system for the  $\Delta \mathbf{v}_i$ , of which (51) is a special case.

Equation (51) poses an alternative: if distance  $\bar{r}$  is to be near the circular light-cone distance  $r_b = \frac{1}{\mu\theta^2}$ , then the last term on the right-hand-side of (51) is negligible and the vanishing of the determinant requires that at least one particle has a *quasi-luminal* velocity. Otherwise collisions can happen only at distances near the very small protonic radius. In variational electrodynamics, 180-degree corners function as *stepping-stones* when linearizations reach the critical quasi-luminal velocities necessary for the formation of corners at-a-far-distance. Above threshold, corner creation is a rebounding mechanism that offers an alternative to charges falling into each other. The stepping-stone condition further implies that the amplitude of transversal mode (50) at layer edge must be  $A \simeq \frac{r_b}{\pi q}$ .

Recalling that motion in the regular layer is almost-transversal, the WECC2 (16) poses another threshold for the 90-degree corner followed by a straight-line quasi-luminal dive (point (a) in FIG. (3)): since  $|\mathbf{v}_1|$  must be continuous, the dive requires a quasi-luminal transversal velocity at layer edge by continuity. Again, the exponential blow-up of Eq. (50) reaches a quasi-luminal velocity at layer edge if  $A \simeq \frac{r_b}{\pi q}$ .

FIG. (4) illustrates the boundary layer motion in the presence of planar and perpendicular perturbations. The green lines of FIG. (4) indicate corners connected by the light-cone condition, again with exaggerated magnitudes to illustrate the gapped-solenoidal motion. The layering must fit an integer number of regular regions and boundary layers in the circle. Assuming the time of flight for each abc segment of FIG. (3) to be  $\Delta t \simeq 2r_b$ , the total time for  $N_c$  corners is  $2\alpha \frac{2\pi}{\Omega} \simeq 2N_c r_b$ , yielding the rough estimates  $\theta \simeq \frac{2\pi\alpha}{N_c} = \frac{(\theta - \epsilon_{xy})}{qN_c}$ .




FIG. 4. Boundary layer motion along a 3-dimensional orbit with perpendicular and transversal perturbations. Perturbed trajectory of (heavier) particle 1 (solid red) and perturbed trajectory 2 (solid blue). Solid green lines indicate corners in light-cone and dashed dark lines are the unperturbed orbits. Boundary layer and delay angle have exaggerated magnitudes for illustrative purposes. Arbitrary units.

## WEIERSTRASS-ERDMANN AND PROBABILITY

The tangent dynamics of circular orbits has an infinite number of *unstable* transversal modes of *arbitrarily large frequencies*, as seen by the non-zero real parts of linear growths (41) and (42). This is unlike the classical Coulombian ODE approximation for the two-body problem, whose tangent dynamics has a finite number of frequencies of the order of the orbital frequency [25]. Since linearized modes (41) and (42) are unstable, continuation along the  $C^\infty$  segment would blow up, and a velocity discontinuity is needed to break from the  $C^\infty$  segment. A corner requires stepping-stone condition (51), and for a *finitely measured* family of orbits with corners to exist, the WECC2 along orbits with planar and perpendicular perturbations requires the resonances studied below.

The perpendicular perturbation in the regular region is constructed analogously to the planar perturbation (50), using the linearized modes explained in appendix B of Ref. [15], and yielding

$$Z_j = B \frac{\cosh(\frac{\sigma_z t}{r_b})}{\cosh(\frac{\sigma_z}{2})} \sin\left(\frac{[\pi s + \epsilon_z(s)](t - \alpha r_b)}{r_b}\right), \quad (52)$$

where  $s$  is any integer (possibly different from  $q$ ) and  $B$  is an arbitrary amplitude smaller than  $\frac{r_b}{\pi s}$  to avoid a superluminal velocity at layer edge. Again, the amplitude of the transversal perturbation (52) is constructed to vanish along the circular orbit at  $t = \alpha r_b$  and  $t = r_b - \alpha r_b$ , to keep the property that corners see other corners in light-cone (a boundary-layer-adjusted resonance).

Since the phase of oscillation ( $\simeq \pi s/r_b$ ) is fast, a large orbital velocity at layer edge results from a small amplitude  $B \simeq \frac{r_b}{\pi s}$ , which is illustrated in FIG. (3). Layer edge

amplitudes of both types of transversal modes should vanish while their derivatives reach a quasi-luminal velocity. Amplitude (50) vanishes at  $t = r_b - \alpha r_b$  when

$$[\pi q + \epsilon_{xy}(q) - \theta](1 - 2\alpha) = \pi q, \quad (53)$$

other multiples of  $\pi$  being excluded because  $\epsilon_{xy}(q)$ ,  $\theta$  and  $\alpha$  are small. Analogously, amplitude (52) vanishes at  $t = r_b - \alpha r_b$  when

$$[\pi s + \epsilon_z(s)](1 - 2\alpha) = \pi s, \quad (54)$$

where again other integer multiples of  $\pi$  are impossible because  $\epsilon_z(s)$  and  $\alpha$  are small.

The climbing/diving edges at  $t = \alpha r_b$  and  $t = r_b - \alpha r_b$  of the *periodic* broken minimizer are equivalent, implying that (50) and (52) with arbitrary amplitudes near  $(A, B) = (\frac{r_b}{\pi q}, \frac{r_b}{\pi s})$  should yield the same quasi-luminal  $|\mathbf{v}_j|$ . For this it is *necessary* that

$$\theta = \frac{q\epsilon_{xy}(q) - s\epsilon_z(s)}{q}. \quad (55)$$

Condition (55) ensures the WECC2 that  $|\mathbf{v}_j|$  is continuous at corners of the periodic broken minimizer and is a *probabilistic argument*. Given (55), at points (a), (c) and (h) of FIG. (3) a *finitely measured* family of orbits include a corner turn, i.e., a *whole neighbourhood* of perturbed orbits near  $(A, B) = (\frac{r_b}{\pi q}, \frac{r_b}{\pi s})$ . Next we calculate the magnitudes of these resonant orbits with  $q = s$ .

For each  $q = s$ , condition (55) determines a unique  $\theta$  together with Eqs. (39) and (40), as listed in Table I. For comparison, Table I also lists the first line of each spectroscopic series, i.e., the circular lines from quantum level level  $k + 1$  to quantum level  $k$ . Historically, the series of hydrogen were named after Lyman, Balmer, Ritz-Paschen, Brackett, etc. The frequency over reduced mass of the first line of each spectroscopic series in atomic units is the second column of Table I. We used a Newton method in the complex- $\lambda$  plane to solve Eqs. (39), (40) and (55), as used in Ref. [15] with Dirac's theory. Our calculations for hydrogen used the protonic-to-electronic mass ratio  $(m_2/m_1) = 1836.1526$ . Table I gives frequency over reduced mass calculated by QM for the first line of the  $k^{th}$  spectroscopic series (atomic units), numerically calculated orbital frequency over reduced mass (for a suitable integer  $q(k)$ ),  $(137^3\Omega)/\mu = 137^3\theta^2(\epsilon_{xy} - \epsilon_z)$ , angular momentum of unperturbed circular guide, and integer  $q$ .

Table I is to be compared with Table I of Ref. [15], which discusses Dirac's electrodynamics with self-interaction. The same surprising agreement is found in Ref. [15], even though Dirac's theory has limitations not shared with the Wheeler-Feynman theory, as discussed in Ref. [2]. Notice we also jumped  $q = 10$  in Ref. [15], again because the resonance condition is only necessary.

| $k$ | $w_{QM}$               | $137^3\theta^2(\epsilon_{xy} - \epsilon_z)$ | $l_z = \theta^{-1}$ | $q$ |
|-----|------------------------|---------------------------------------------|---------------------|-----|
| 1   | $3.750 \times 10^{-1}$ | $3.852 \times 10^{-1}$                      | 188.32              | 7   |
| 2   | $6.944 \times 10^{-2}$ | $8.919 \times 10^{-2}$                      | 306.67              | 9   |
| 3   | $2.430 \times 10^{-2}$ | $2.485 \times 10^{-2}$                      | 469.54              | 11  |
| 4   | $1.125 \times 10^{-2}$ | $1.392 \times 10^{-2}$                      | 569.61              | 12  |
| 5   | $6.111 \times 10^{-3}$ | $8.070 \times 10^{-3}$                      | 683.13              | 13  |
| 6   | $3.685 \times 10^{-3}$ | $4.825 \times 10^{-3}$                      | 810.89              | 14  |
| 7   | $2.406 \times 10^{-3}$ | $2.966 \times 10^{-3}$                      | 953.65              | 15  |
| 8   | $1.640 \times 10^{-3}$ | $1.870 \times 10^{-3}$                      | 1112.21             | 16  |
| 9   | $1.173 \times 10^{-3}$ | $1.206 \times 10^{-3}$                      | 1287.27             | 17  |
| 10  | $8.678 \times 10^{-4}$ | $7.942 \times 10^{-4}$                      | 1479.62             | 18  |
| 11  | $6.600 \times 10^{-4}$ | $5.329 \times 10^{-4}$                      | 1690.02             | 19  |
| 12  | $5.136 \times 10^{-4}$ | $3.639 \times 10^{-4}$                      | 1919.20             | 20  |

TABLE I. Numerical calculations for hydrogen with  $(m_2/m_1) = 1836.1526$ . Quantum number  $k$  of the circular Bohr transition  $k+1 \rightarrow k$ , frequency over reduced mass of the circular QM line in atomic units,  $w_{QM} \equiv \frac{1}{2}(\frac{1}{k^2} - \frac{1}{(k+1)^2})$ , orbital frequency over reduced mass in atomic units,  $(\Omega/\mu) = 137^3\theta^2(\epsilon_{xy} - \epsilon_z)$ , angular momentum of unperturbed orbit in units of  $e^2/c$ ,  $l_z = \theta^{-1}$ , and integer  $q$ .

As mentioned below Eqs. (48) and (49), the energetic mismatches are monotonically increasing for  $q \geq 6$ , and Table I frequencies are within ten percent agreement with the first 12 circular hydrogen lines starting from  $q = 7$  at  $k = 1$ , i.e.,  $q(1) = 7$ . Since condition (55) is only *necessary* (and not sufficient), some values of  $q$  may correspond to unstable orbits. This is analogous to the description by QM [9], where there are selection rules on top of *three* conditions involving integer quantum numbers, and Table II includes the lines that were skipped in Table I. A theory for the  $q$ 's that were skipped is presently lacking. Inspection of Table II shows that for  $q = 1, 2, 3, 4, 5, 6$  condition (55) predicts  $\theta$  still in the atomic range, but the angular momentum spacing is about half of Planck's constant. Table II also includes the skipped numerical calculations for  $q = 8$  and  $q = 10$ . A suggestion from Sommerfeld's quantization is that there must be *three* conditions like (55), involving *three* integer quantum indices, not just one, reinforcing that our single condition (55) is only necessary and alone might not determine a stable orbit. Inspection shows that if the values of Table II were included in Table I, the angular momentum jump from consecutive lines would be much lesser than about a hundred units of  $e^2/c$ , which is suggestive of what the missing conditions should do.

Next we show the numerical calculations for the muonium atom using the positive-muon-to-electron mass ratio  $(m_2/m_1) = 1836.1526/9$ . Table III lists the frequency over reduced mass of the first line of each spectroscopic series as calculated with QM (in atomic units), orbital frequency in atomic units,  $(137^3\Omega)/\mu = 137^3\theta^2(\epsilon_{xy} - \epsilon_z)$ , and angular momentum of unperturbed circular orbit in units of  $e^2/c$ . The agreement of the numerical calculations with nature and QM is again within a few percent

| $137^3\theta^2(\epsilon_{xy} - \epsilon_z)$ | $l_z = \theta^{-1}$ | $q$ |
|---------------------------------------------|---------------------|-----|
| 1.506                                       | 119.53              | 1   |
| 10.653                                      | 62.27               | 2   |
| 9.466                                       | 64.77               | 3   |
| 4.664                                       | 82.01               | 4   |
| 2.006                                       | 108.63              | 5   |
| $8.622 \times 10^{-1}$                      | 143.96              | 6   |
| $1.808 \times 10^{-1}$                      | 242.32              | 8   |
| $4.6095 \times 10^{-1}$                     | 382.16              | 10  |

TABLE II. Numerical calculations for hydrogen with  $q < 7$  and  $(m_2/m_1) = 1836.1526$ . orbital frequency over reduced mass in atomic units,  $(\Omega/\mu) = 137^3\theta^2(\epsilon_{xy} - \epsilon_z)$ , angular momentum of unperturbed circular guide in units of  $e^2/c$ ,  $l_z = \theta^{-1}$ , and integer  $q$ .

for frequencies.

| $k$ | $w_{QM}$               | $137^3\theta^2(\epsilon_{xy} - \epsilon_z)$ | $l_z = \theta^{-1}$ | $q$ |
|-----|------------------------|---------------------------------------------|---------------------|-----|
| 1   | $3.750 \times 10^{-1}$ | $4.039 \times 10^{-1}$                      | 185.63              | 7   |
| 2   | $6.944 \times 10^{-2}$ | $8.762 \times 10^{-2}$                      | 308.94              | 9   |
| 3   | $2.430 \times 10^{-2}$ | $2.356 \times 10^{-2}$                      | 478.65              | 11  |
| 4   | $1.125 \times 10^{-2}$ | $1.304 \times 10^{-2}$                      | 582.96              | 12  |
| 5   | $6.111 \times 10^{-3}$ | $7.491 \times 10^{-3}$                      | 701.31              | 13  |
| 6   | $3.685 \times 10^{-3}$ | $4.445 \times 10^{-3}$                      | 834.53              | 14  |
| 7   | $2.406 \times 10^{-3}$ | $2.717 \times 10^{-3}$                      | 983.41              | 15  |
| 8   | $1.640 \times 10^{-3}$ | $1.704 \times 10^{-3}$                      | 1148.75             | 16  |
| 9   | $1.173 \times 10^{-3}$ | $1.095 \times 10^{-3}$                      | 1331.34             | 17  |
| 10  | $8.678 \times 10^{-4}$ | $7.187 \times 10^{-4}$                      | 1531.19             | 18  |
| 11  | $6.600 \times 10^{-4}$ | $4.810 \times 10^{-4}$                      | 1751.36             | 19  |
| 12  | $5.136 \times 10^{-4}$ | $3.277 \times 10^{-4}$                      | 1990.34             | 20  |

TABLE III. Numerical calculations for muonium with  $(m_2/m_1) = 1836.1526/9$ . Quantum number  $k$  of the circular Bohr transition  $k+1 \rightarrow k$ , frequency over reduced mass of the circular QM line in atomic units,  $w_{QM} \equiv \frac{1}{2}(\frac{1}{k^2} - \frac{1}{(k+1)^2})$ , orbital frequency over reduced mass in atomic units,  $(\Omega/\mu) = 137^3\theta^2(\epsilon_{xy} - \epsilon_z)$ , angular momentum of unperturbed circular guide in units of  $e^2/c$ ,  $l_z = \theta^{-1}$ , and integer  $q$ .

We repeat the numerical calculations for positronium using the positron-to-electron mass ratio  $(m_2/m_1) = 1$ : Table IV lists the frequency over reduced mass of the first line of each spectroscopic series calculated by QM (in atomic units), orbital frequency over reduced mass in atomic units and the angular momentum of the unperturbed circular orbit in units of  $e^2/c$ . Notice in Table IV that for positronium the values of  $l_z = 1/\theta$  are consistently larger. Using again the fact that condition (55) is only necessary, Table IV starts the  $q$  when the angular momentum spacing is about constant, i.e.,  $q = 8$ . For positronium the numerical calculations find the first root  $1/\theta = 40.501$  only at  $q = 3$ , *again* in the atomic magnitude. The spectrum agrees with QM within less than a few percent for the circular lines of the first 12 series.

In the days of Bohr, only about 12 lines of the Balmer series could be observed with vacuum tubes, and about

| $k$ | $w_{QM}$               | $137^3\theta^2(\epsilon_{xy} - \epsilon_z)$ | $l_z = \theta^{-1}$ | $q$ |
|-----|------------------------|---------------------------------------------|---------------------|-----|
| 1   | $3.750 \times 10^{-1}$ | $3.423 \times 10^{-1}$                      | 246.76              | 8   |
| 2   | $6.944 \times 10^{-2}$ | $7.749 \times 10^{-2}$                      | 404.86              | 10  |
| 3   | $2.430 \times 10^{-2}$ | $2.189 \times 10^{-2}$                      | 617.037             | 12  |
| 4   | $1.125 \times 10^{-2}$ | $1.240 \times 10^{-2}$                      | 745.61              | 13  |
| 5   | $6.111 \times 10^{-3}$ | $7.286 \times 10^{-3}$                      | 890.33              | 14  |
| 6   | $3.685 \times 10^{-3}$ | $4.416 \times 10^{-3}$                      | 1052.06             | 15  |
| 7   | $2.406 \times 10^{-3}$ | $2.753 \times 10^{-3}$                      | 1231.64             | 16  |
| 8   | $1.640 \times 10^{-3}$ | $1.759 \times 10^{-3}$                      | 1429.92             | 17  |
| 9   | $1.173 \times 10^{-3}$ | $1.149 \times 10^{-3}$                      | 1647.73             | 18  |
| 10  | $8.678 \times 10^{-4}$ | $7.667 \times 10^{-4}$                      | 1885.88             | 19  |
| 11  | $6.600 \times 10^{-4}$ | $5.209 \times 10^{-4}$                      | 2145.20             | 20  |
| 12  | $5.136 \times 10^{-4}$ | $3.599 \times 10^{-4}$                      | 2426.50             | 21  |

TABLE IV. Numerical calculations for positronium with  $(m_2/m_1) = 1$ . Quantum number  $k$  of the circular Bohr transition  $k+1 \rightarrow k$ , frequency over reduced mass of the circular QM line in atomic units,  $w_{QM} \equiv \frac{1}{2}(\frac{1}{k^2} - \frac{1}{(k+1)^2})$ , orbital frequency over reduced mass in atomic units,  $(\Omega/\mu) = 137^3\theta^2(\epsilon_{xy} - \epsilon_z)$ , angular momentum of unperturbed orbit in units of  $e^2/c$ ,  $l_z = \theta^{-1}$ , and integer  $q$ .

thirty-three from spectra of celestial bodies [26]. Surprisingly, the emission frequencies agree better with QM for the decays from the 12 deepest quantum levels. As explained in [15], the cancelation of dipolar far-fields involves quadratic terms that might require larger amplitudes at large  $q$ . Given that the  $xy$  modes modify the unperturbed  $z$ -angular momentum more and more at larger  $q$ , our perturbative results should get worse at larger  $q$ . We carried our numerical calculations up to  $q = 43$  (not shown in Table II), and observed that the angular momentum spacing increases slowly with  $q$ . By defining an effective  $k \equiv (l_z/170)$ , the agreement of the numerical calculations with an effective angular momentum separation seems to continue indefinitely within thirty percent, suggesting we can approximate a large number of eigenvalues near the discrete spectrum of Schroedinger's equation. The agreement with emission lines above  $k = 13$  slowly deteriorates, which is suggestive that the corresponding broken minimizers are becoming far from planar. A one-to-one comparison with natural spectra should wait the investigation of broken extrema with spikes filling a tridimensional region.

The surprising agreement of the numerical calculations with a universal value for the fine-structure constant is due to the logarithmic dependence of  $\sigma \equiv \ln(\sqrt{\frac{4M}{\mu\theta^4}})$ , as explained above Eq. (37). Moreover, because of the weak dependence on  $\theta$ , Eq. (56) is an implicit transcendental condition and the  $O(1/\lambda^2)$  formulas (48) and (49) yield

$$\frac{1}{\theta} = \frac{1}{\epsilon_{xy} - \epsilon_z} = \frac{(\sigma^2 + \pi^2 q^2)^2}{8\pi\sigma q}. \quad (56)$$

The roots of (56) are insensitive to changes in  $(\mu/M)$  over three orders of magnitude, e.g., for  $q = 7$  and  $(\mu/M) = 1/1837$  (hydrogen), Eq. (56) yields  $\theta \simeq 190.09$ , while

for  $q = 7$  and  $(\mu/M) = 9/1836$  (muonium), Eq. (56) yields  $\theta \simeq 187.17$ , and last for  $q = 7$  and  $(\mu/M) = 1/4$  (positronium), Eq. (56) yields  $\theta \simeq 186.81$ . The numerically calculated values of  $l_z \equiv \theta^{-1}$  fall approximately between the two consecutive Bohr orbits  $k$  and  $k+1$  (artificially) used to calculate frequencies, and Tables I, III and IV show that frequencies of spectral lines agree even better with QM and nature.

Last, the orbital frequencies determined by the Weierstrass-Erdmann mechanism (55) can be expressed as a difference of two spectroscopic terms, just like in QM, i.e.,

$$\Omega \equiv \mu\theta^3 = \frac{\theta\epsilon_{xy}(q) - \theta\epsilon_z(s)}{(1/\theta)}, \quad (57)$$

with spectroscopic terms  $\theta\epsilon_{xy}(q)$  and  $\theta\epsilon_z(s)$  defined by the eigenvalues of two *linear* and *infinite dimensional* problems, i.e., Eq. (31) and Eq. (B16) of Appendix B in Ref. [15].

## GAC AND VANISHING FAR-FIELDS

Radiation in Maxwell's electrodynamics is treated with Poynting's theorem [5], which within variational electrodynamics holds only for extended fields of globally  $C^2$  orbits, in which case Maxwell's electrodynamics is a special case of variational electrodynamics. In the following we use only generally valid properties of variational electrodynamics. The physical meaning of the extended far-fields in variational electrodynamics is only that of a coupling to distant charges. The significance of a surface integral of the Poynting vector of extended fields undefined in sets of zero measure is an open application of the divergence theorem of modern functional analysis [10].

Natural spectra of atomic gases containing Avogadro's number of atoms pose a many body problem where each atom suffers perturbations from all other atoms. This unless a decoupling from other atoms or laboratory boundaries is achieved by the GAC. In Ref. [2] it was shown that globally bounded two-body orbits with vanishing extended far-fields *must* involve velocity discontinuities along the continuous trajectories.

Under the above physical interpretation of extended far-fields as couplings, we recall the far-field-vanishing mechanism of Ref. [15] using the quadratic correction to far-field (4) due to the last right-hand-side term of (5), i.e.,

$$\delta\mathbf{E}_{j\pm}^{(2)} \equiv \pm q_j \frac{\mathbf{n}_{j\pm} \times (\mathbf{v}_{j\pm} \times \mathbf{a}_{j\pm})}{r_{j\pm}}. \quad (58)$$

Not surprisingly, quadratic term (58) is ignored when approximating the two-body motion with a Coulombian ODE. Again, Colombian ODE's have a tangent dynamics spanned by a finite number of slow frequencies [25],

which cause small accelerations, while the tangent dynamics defined by NDDE (31) has *arbitrarily large* frequencies *and* arbitrarily large accelerations. Since accelerations are arbitrarily large and transversal velocities quasi-luminal, the quadratic term (58) introduces beats of the fast frequencies [15]. The GAC mechanism of Ref. [15] to cancel the dipolar far-fields of the unperturbed (slow) circular motion by (58) with a resonant beat among fast planar and fast perpendicular modes would be inexistent for a Coulombian ODE.

It is a nice surprise that the WECC are compatible with the GAC. In fact, (55) with  $q = s$  is equivalent to the condition for the quadratic term to cancel the dipolar far-fields of the unperturbed circular orbit [15]. Equation (55) with  $q = s$  is exactly Eq. (56) of Ref. [15]. Therefore, (55) determines the finitely measured family of orbits that *can* acquire vanishing far-fields by linear perturbations, as necessary to decouple from experimental boundaries or other atoms of the gas.

The logical application of multiscale perturbation is to balance the fast timescale *first*. For that one could freeze the slow guiding motion more generally than as a circular orbit. As long as the slow velocity is of the order determined by the centripetal force of the Coulombian limit, i.e.,  $\mu v^2 \simeq 1/r$ , the far-field mechanism of  $O(1/\lambda^2)$  corrections to (33) and (36) prescribes the same resonance condition (55). This is further indication that our determination of atomic scales is not accidental, and other finitely measured families of orbits with vanishing far-fields should be found again in the atomic magnitude.

A second confirmation that the WECC mechanism selects the atomic scale is that the simple 180-degree rebounding mechanism turns *off* at smaller scales. According to Eq. (51), 180-degree corners are an alternative to opposite charges falling into each other only if  $\bar{r} \geq \frac{1}{\sqrt{m_1 m_2}}$ , at which distance the mechanism ceases to operate, thus creating a *confinement core* at the *nuclear magnitude*  $\bar{r} < \frac{1}{\sqrt{m_1 m_2}}$ .

We do not claim to have found all possible spiky orbits, and the  $q = s$  family of periodic orbits with vanishing far-fields precisely at the atomic magnitude suggests a systematic search should continue. For non-harmonic (e.g. elliptical) orbits, an effective mechanism of cancelation of the main harmonic of the slow periodic guide should operate for  $q \neq s$ . Investigations of stability are better done numerically, and for that finite element methods are being developed [18, 23].

## DISCUSSIONS AND CONCLUSION

Here, as in [18], we adopted *particle time* for each particle's evolution parameter. The Fokker action with two arbitrary evolution parameters [1], is *improper* for the finite-element method of [18] because the parametrization-invariance symmetry causes a zero

eigenvalue and a singular Hessian matrix [18]. WECC from the parametrization-invariant action were used in [27] to model double-slit interference caused by interaction-at-a-distance with the corner points of the trajectories of material electrons inside the grating [27]. The parametrization-invariant action yields four scalar conditions for the WECC1, while WECC2 is trivially satisfied [28]. The choice of particle-time evolution parameter yields *three* scalar conditions for the WECC1 and *one* scalar condition for WECC2 (an equivalent set of four conditions as explained in [28]).

The GAC uncoupling mechanism operates even at closer distances along globally bounded trajectories of atoms with a zero net charge, because then also the near fields vanish as a consequence of Gauss's integral law. Along trajectories satisfying the GAC, atoms with a zero net charge can stand decoupled at distances ranging from intermediate to far distances. The minimizer for the combined system is the direct product of the unperturbed trajectories precisely because each atom produces no far fields and little near fields to act back on the other atoms.

A motivation for Wheeler-Feynman theory [6] was Sommerfeld's quantization conditions [20, 21, 29]. Surprisingly, broken extrema of variational electrodynamics satisfy a condition involving *precisely* the same relevant quantities of quantum mechanics, i.e., *momenta* and *energies* [4]. Energy and momentum entered Sommerfeld's theory via Hamiltonian mechanics, which became paradoxical after the no-interaction theorem proved that only trivial non-interactive two-body motion is covariantly described by a Hamiltonian [16, 30, 31].

Variational electrodynamics is also free of the Aharonov-Bohm paradox [32] along broken extrema. The origin of this paradox is that along  $C^\infty$  orbits the electromagnetic equations of motion involve only *derivatives* of the vector potential: The Euler-Lagrange equation of partial Lagrangian (15) yields (1) with a right-hand-side equal to  $\nabla U_j + \partial \mathbf{A}_j / \partial t - \mathbf{v}_i \times (\nabla \times \mathbf{A}_j)$ . Instead, along *broken extrema* the vector potential  $\mathbf{A}_j$  *itself* appears on the WECC1, Eq. (14), determining an interference-at-a-distance just like in QM, as discussed in [27].

There is no fitting parameter in our calculations, and the agreement with nature proves variational electrodynamics is a physically sound extension. Agreement beyond thirty percent should wait a perturbative scheme with corrections for the guiding center motion.

The general hydrogenoid spectrum falls in the class  $\Omega_{QM} \equiv \frac{\mu}{2}(\frac{1}{k^2} - \frac{1}{j^2})$  [9]. If variational electrodynamics is to approximate the whole hydrogenoid spectrum, broken minimizers still to be discovered should correspond to the elliptical lines ( $j \neq k+1$ ). For example, spiky orbits with *two* guiding circles, with each particle diving to a lower circle and riding along it for a while, then jumping back to an upper circle and riding along it for a while, and so on. These orbits involve *two* Weierstrass-Erdmann conditions and *two* integers, just like the hydrogenoid series

[9, 26]. A classification of spiky orbits and a multiscale scheme starting from fast motions about non-smooth and almost everywhere slow guiding motions is lacking.

## ACKNOWLEDGEMENTS

Author acknowledges the partial support of a FAPESP regular grant, Marcel Novaes for reading the manuscript and Daniel Câmara for discussions.

---

\* [deluca@ufscar.br](mailto:deluca@ufscar.br)

- [1] J. De Luca, *J. Math. Phys.* **50**, 062701(2009).
- [2] J. De Luca, *Physical Review E* **82**, 026212 (2010).
- [3] John L. Troutman, *Variational Calculus and Optimal Control: Optimization with Elementary Convexity*, second edition, Springer (1995).
- [4] I. M. Gelfand and S. V. Fomin, *Calculus of Variations*, Dover (2000).
- [5] J. D. Jackson, *Classical Electrodynamics* Second Edition, John Wiley and Sons, New York (1975).
- [6] J. A. Wheeler and R. P. Feynman, *Rev. Mod. Phys.* **17**, 157 (1945); J. A. Wheeler and R. P. Feynman, *Rev. Mod. Phys.* **21**, 425 (1949).
- [7] J. Hale, *Theory of Functional Differential Equations*, Springer-Verlag (1977).
- [8] A. Bellen and M. Zennaro, *Numerical Methods for Delay Differential Equations* Oxford University Press, NY (2003).
- [9] H. A. Bethe and E. E. Salpeter, *Quantum Mechanics of One- and Two-Electron Atoms*, Dover Ed. (2008), ISBN. 0-486-46667-1
- [10] H. Brezis, *Functional Analysis, Sobolev Spaces and Partial Differential Equations*, Springer, New York (2010).
- [11] A. Schild, *Phys. Rev.* **131**, 2762 (1963); A. Schild, *Science* **138**, 994 (1962).
- [12] R.D. Driver, *Physical Review D* **19**, 1098 (1979).
- [13] A. Staruszkiewicz, *Acta Physica Polonica XXXIII*, 1007 (1968).
- [14] C.M. Andersen and H.C. von Baeyer, *Phys. Rev. D* **5**, 2470 (1972).
- [15] J. De Luca, *Physical Review E* **73**, 026221 (2006).
- [16] *The Theory of Action-at-a-Distance in Relativistic Particle Dynamics*, edited by E. H. Kerner, Gordon and Breach, New York, 1972.
- [17] F. Hoyle and J. Narlikar, *Cosmology and Action at a Distance Electrodynamics*, World Scientific Publishing, Singapore (1996).
- [18] J. De Luca, A. R. Humphries and S. B. Rodrigues, *Journal of Computational and Applied Mathematics* **236**, 3319-3337 (2012).
- [19] J. L. Kaplan, M. Sorg, and J.A. Yorke, *Nonlinear Analysis, Theory, Methods and Applications* **3**, 53 (1979).
- [20] C. M. Andersen and Hans C. von Baeyer, *Annals of Physics* **60**, 67-84 (1970).
- [21] H. C. von Baeyer, *Physical Review D* **12**, 3086 (1975).
- [22] D. Leiter, *Am. J. Phys.* **38**, 207 (1970).
- [23] Y. Jabri, *The mountain pass theorem*, Cambridge University Press, ISBN 0 521 82721 3, Cambridge (2003).
- [24] P. A. M. Dirac, *Proceedings of the Royal Society of London, ser. A* **167**, 148 (1938).
- [25] J. De Luca, *Phys. Rev. Lett.* **80**, 680 (1998), J. De Luca *Phys. Rev. E* **58**, 5727 (1998).
- [26] N. Bohr, *Philos. Mag.* **26**, 1 (1913); **26**, 476 (1913).
- [27] J. De Luca, *Journal of Computational and Theoretical Nanoscience* **8**, 1040-1051 (2011).
- [28] A. O. Barut, *Electrodynamics and Classical Theory of Fields and Particles*, Dover (NY), (1980).
- [29] Sommerfeld, Arnold (1919) *Atombau und Spektrallinien*. Braunschweig: Friedrich Vieweg und Sohn. ISBN 3-87144-484-7 and ter Haar, D. *The Old Quantum Theory*. Pergamon Press (1967). ISBN 0-08-012101-2.
- [30] D. G. Currie, T. F. Jordan, and E. C. G. Sudarshan, *Rev. Mod. Phys.* **35**, 350 (1963).
- [31] G. Marmo, N. Mukunda, and E. C. G. Sudarshan, *Phys. Rev. D* **30**, 2110 (1984).
- [32] Y. Aharonov and D. Bohm, *Physical Review* **115**, 485 (1959).